15 tests: FAirEST 15

Circle the most accurate answer for each assessment

<b>1. Facial Pattern</b> A. Mesofacial	B. Brachyfacial (wide)	C. Dolichofacial (long)	
<b>2. Facial Profile</b> A. Normal	B. Convex	C. Concave	
<b>3. Dental Profile</b> A. Class I	B. Class II	C. Class III	
<b>4. Tongue Range of Motion Rat</b> A. Grade 1 (>80%)	<b>tio (TRMR)</b> B. Grade 2 (50-80%)	C. Grade 3 (<50%)	D. Grade 4 (<25%)
<b>5. Kotlow Tongue Tie</b> A. Normal or mild (12-16mm)	B. Moderate (8-12mm)	C. Severe (4-8mm)	D. Complete (0-4mm)
<b>6. Kotlow Lip Tie Attachment:</b> A. Mucosal	B. Gingival	C. Interdental papilla	D. Palatine papilla
<b>7. Mentalis Strain</b> A. No chin strain present	B. Mild chin strain	C. Moderate chin strain	D. Severe chin strain
<b>8. Mallampati</b> A. Class 1	B. Class 2	C. Class 3	D. Class 4
9. Tonsil size (Brodsky scale) A. Grade 1 (<25%)	B. Grade 2 (26-50%)	C. Grade 3 (51-75%)	D. Grade 4 (>75%)
10. Maxillary Intercanine DistanceA. Wide (>37mm)B. Neutral (31-37mm)C. Narrow (<31mm)			
<b>11. Maxillary Intermolar Distan</b> A. Wide (>52mm)	n <b>ce (Mesiobuccal cusp)</b> B. Neutral (46-52mm)	C. Narrow (<46mm)	
<b>12. Incisor display @ Rest</b> A. 0-25%	B. 26-50%	C. 51-75%	D. 76-100%
<b>13. Bruxism</b> A. No wear detected	B. Mild dental wear	C. Moderate dental wear	D. Severe dental wear
<b>14. Swallowing Tongue Thrust</b> A. Swallow with ease	Compensation Test B. Swallow with difficulty	C. Unable to swallow without tongue thrust compensation	
<b>15. Nasal Breathing Test</b> A. 3+ min	B. 2-3 min	C. 1-2 min	D. <1 min
2 of 2 Complete both sides			

www.fairest.org © Copyright 2019-2020 – Free to copy and use without alterations for non-commercial use

Functional Airway Evaluation Screening Tool

15 tests: FAirEST 15

	1	2	3	4
Nasal vs. Mouth Breathing	Primarily nasal breathing: Rarely to never mouth breathes when awake.	Sometimes mouth breathes when awake.	Often mouth breathes when awake.	Almost always mouth breathes when awake.
	Primarily nasal breathing: Rarely to never mouth breathes when asleep.	Sometimes mouth breathes when asleep.	Often mouth breathes when asleep.	Almost always mouth breathes when asleep.
Posture	Rarely slouches.	Sometimes slouches.	Often slouches.	Almost always slouches.
	Primarily sleeps on back.	Sleeps on back > more than sides or stomach.	Sleeps on sides/stomach > more than back.	Primarily sleeps on sides or stomach.
CNS	Rarely to never has difficulty concentration. Easy to focus.	Sometimes has difficulty concentrating. Somewhat difficult to focus.	Often has difficulty concentrating. Moderately difficult to focus.	Almost always has difficulty concentrating. Very difficult to focus.
	Almost always relaxed. Rarely feels anxious.	Often relaxed. Sometimes feels anxious.	Often stressed. Sometimes feels relaxed.	Almost always stressed. Rarely feels relaxed.

## Check all the following that apply:

- \_\_\_\_Prior tongue-tie release
- \_\_\_\_Prior tonsillectomy and/or adenoidectomy
- \_\_\_\_Prior orthodontics
- \_\_\_\_Prior jaw or maxillofacial surgery
- \_\_\_\_Prior treatment with myofunctional therapy
- \_\_\_\_Craniofacial anomaly, cerebral palsy, or syndromic facies
- \_\_\_\_None of the above apply

1 of 2 Complete both sides